Deer carcasses, scavengers, soil nutrients and invertebrates

Scott Newey (scott.newey@hutton.ac.uk)

Justin Irvine, Debbie Fielding, and Rene van der Wal



The James Hutton Institute

www.hutton.ac.uk/deerdecomposition





- Introduction, background and why?
- Studies and approaches used
- (Preliminary) Results
 - Vertebrate Scavengers
 - Soil Nutrients
 - Invertebrates
- Summary

Introduction: Red deer





"Iconic" species




"Pest" species

- Culturally, economically and ecologically important
- National herd ~ 360,000 400,000



Sport shooting & culling mortality ~ 15%





Mortality, biodiversity, culls and venison




Natural mortality ~ 1%



Carrion and carcass provisioning





 Reduced amount of carrion in the landscape loss of nutrients and energy break down in nutrient cycling loss of fertility decline in scavengers



 Carcass provisioning restore ecosystem processes as a conservation tool

Fielding, Newey, Irvine & van der Wal (2014) AMBIO 43(6) 810-891

Carcass provisioning and controversies



38 DEGREES

Q Search

Start a Campaign News

STOP THE JOHN MUIR TRUST FROM MASSACRING OUR SCOTTISH RED DEER.



TO: PAUL MONAGHAN MP

Please stop the John Muir Trust from killing any more Red Deer in Assynt and Knoydarl.

Why is this important?

Red deer are native to our wild lands and one of the many reasons why tourists and cash rich stalkers come to valit these generally economically disadvantaged areas. The Red Deer are beautiful majestic animals and must be managed in a way as to assure there is a viable population for all JMT preach "deer management" which to them means kill on sight to

000

NEWS.

Fresh clash in Highland deer cull row

Transvery 2011 The Calves Book

10.0



The MT said Bynaserds of door de a paintal dooth of starvation avery yea hillsides because of overpopulation - and that the particle spins were tryin "deflect attention" from sorutiny as Sociated penders greater land reform.

Gamekeepers angered by John Muir Trust deer cull



The SGA is calling on the Scottish Government to investigate claims that the john Muir Drust left.



Call to investigate Highland deer cull after dozens of carcusses are left to rot on hill

00000

23.5 member 2015, by Colum Auss

10 guardenes




The Scottah Government has been urged to investigate claims a conservation charity left dozens of deer to not on a Highland hilblde.



Effects of deer carcasses on upland food webs

Three complimentary but independent questions:

- 1. Do, and if so what, scavengers use deer carcasses?
- 2. What are the spatial and temporal effects?
- 3. How does land management influence carcass decomposition?





Approaches and studies





- Motivations for carcass provisioning
 - Fielding *et al*. (2014) AMBIO 43(6) 810-891
- Cross-sectional study
- Intensive carcass placement study
- Extensive carcass placement study
 - vertebrate scavengers
 - invertebrates
 - vegetation,
 - soil microbes
 - soil fungi (mycorrhizae)
 - soil nematodes
 - soil nutrients

Extensive carcass placement study

- Six study sites; 2013-2015
- Different land management objectives











Animal species using carcasses

Species/Management	Legal predator control	No predator control
Crow/Raven	+	+
Buzzard	-	+
Short-eared owl	+	+
Golden eagle	-	+
Fox	+	+
Pine marten	-	+
Stoat/weasel	-	+
Badger	-	+
Otter	-	+
Cat (unknown sp.)	-	+



The James Hutton Institute













cted & CAMPIE ANTIC

03-18-2013 12:40 #i

Effects of land management


3-4 months after placement



No predator control

Predator control








The James Hutton Institute

Invertebrates: Silphidia







Summary and conclusions



- There is national and international interest in carcass^{In} provisioning as a rewilding and conservation tool.
- Controversial and benefits and disbenefits have not been quantified.
- A range of vertebrate scavenger species visited carcasses, though population level effects remain unknown.
- Land management effected the species detected and the decomposition of carcasses.
- Carcasses appear to have a large and spatially limited effect on soil nutrients, and invertebrate communities and abundance.



Thank s to all those who have helped during the project, and to all the estates that provided access and assisted with this study. Funded by



The James Hutton Institute is supported by the Scottish Government's Rural and Environment Science and Analytical Services Division (RESAS)

www.hutton.ac.uk/deerdecomposition